

THURSDAY

19th MAY

Eliseo Theatre, via Roma 73 - Nuoro

ROOM_A

TOPIC 1

Agroforestry and the environment

T1.1 - Climate change (adaptation & mitigation)

Parallel session co-hosted by the project SALAM-MED

Chair - Pier Paolo Roggero, University of Sassari, Department of Agricultural Sciences

- 09:00** **O_1.1_21 Sustainable approaches to land and water management in Mediterranean drylands: the SALAM-MED project**
Chiara Ceseracciu, Elena Bresci, Mauro Centritto, Nicola Lamaddalena, Ehab Zaghoul, Mongi Ben Zaied, Stavros Solomos, Zein Kallas, Said Wahbi, Antonio Del Campo, Houssemedin Henchir, Alessio Merella, Andrea Galante, Jean Marc Fures, Youssef Brahimi, Taysir Arbasi, Pier Paolo Roggero
- 09:15** **O_1.1_7 DIGITAF, a European project to help agroforestry meet climate, biodiversity and farming sustainability goals: linking field and cloud**
Marie Gosme, Bert Reubens, Gerry Lawson, Paul J. Burgess, Sonja Kay, Marco DeBoer, Patrick Worms
- 09:30** **O_1.1_19 Harvest index and nitrogen uptake of barley in intercropped walnut orchard**
Helena Žalac, Ante Bubalo, Vladimir Ivezić, Jurica Jović, Miro Stošić, Vladimir Zebec
- 09:45** **O_1.1_20 Framework for building climate change resilience into landscape restoration through agroforestry: an England case study**
Will Simonson, Colin Tosh, David Wolfe, Daan Verstand, Saskia Houben, Isabella Selin Norén, Michelle Allen

Harvest index and nitrogen uptake of barley in intercropped walnut orchard

EURAF 2022
Agroforestry for the Green Deal transition.
Research and innovation towards the
sustainable development of agriculture and
forestry

Corresponding Author:
vivezic@fazos.hr

Helena Žalac¹, Ante Bubalo¹, Vladimir Ivezić¹, Jurica Jović¹, Miro Stošić¹, Vladimir Zebec¹

¹ University of J.J. Strossmayer in Osijek, Faculty of Agriobiotechnical Sciences Osijek, Vladimira Preloga 1, 31000 Osijek, Croatia (e-mail: vivezic@fazos.hr)

Theme: Climate Change (adaptation & mitigation)

Keywords: barley, intercropping, harvest index, nitrogen use efficiency

Abstract

Introduction. Intercropping of annual crops with woody species has shown to be a great ecologically sustainable alternative to conventional agricultural systems. Besides ecological benefits, intercropped systems may be more efficient in terms of productivity and resource capture. Trees can help intercrops acquire nutrients by different mechanisms; absorption of nutrients from deeper soil layers outside the root zone of agricultural crops or absorption of nutrients in chemical forms that are not available for crops, and fixation of atmospheric nitrogen (Cannell et al. 1996). Aim of our research is to investigate barley productivity and nitrogen use efficiency in an intercropped system with common walnut (*Juglans regia*). **Materials and methods.** The trial was set up in Eastern Croatia near city of Đakovo. It consisted of a barley monoculture plot and an intercropped walnut orchard. The orchard was 12 years old with a distance between tree rows of 8 m and barley was sown in strips of 6 m width from October 2019 until June 2020. At harvest, plant samples were collected from 0,5 m² subplots where the plants were cut at the soil level and grain and straw were weighted separately for determination of yield and harvest index. Harvest index was calculated as ratio of barley grain yield and total aboveground biomass. From dried samples, nitrogen concentration was determined by Kjeldahl digestion method. Nitrogen uptake (whole plant) and nitrogen use efficiency (NUE) were calculated as follows:

$$\text{N-uptake (kg ha}^{-1}\text{)} = \text{N\%} * \text{DM}/100 \quad \text{eq. 1}$$

$$\text{NUE (kg kg}^{-1}\text{)} = \text{Grain Yield}/\text{Nuptake} \quad \text{eq. 2}$$

Results. Barley yield in monoculture plot was 7383.86 kg ha⁻¹ and in intercropped orchard 7094.59 kg ha⁻¹ (per cropped area). Since intercropped orchard had generally better soil properties than monoculture plot, this yield reduction is probably due to shading effect by walnut trees. However, harvest index (HI) was significantly greater in intercropped orchard and accordingly had significantly lower total aboveground biomass. This observation suggests that barley in intercropped orchard allocated more assimilates to grains in expense of biomass production. Since HI strongly depends on water availability, this improved reproductive/yield efficiency could be a result of improved microclimatic conditions in intercropped orchards (Gosme et al., 2016). Furthermore, strong correlation was found between HI and NUE ($r=0.81$, $p<0.01$). Although barley in monoculture plot has taken up more N in kg ha⁻¹, intercropped barley had better NUE in terms of producing more grain yield per N taken up. Also, intercropped barley had higher grain N content than monoculture barley, which is a result of better N remobilization. This high N content, however, is negatively correlated with the final grain yield due to the dilution effect. This observation could be connected with shading by walnut trees and it is in accordance with results obtained by Artru et al. (2016) and Dufour et al. (2012) who found the highest grain protein content in wheat grown in continuous/uniform shade. In conclusion, our results show that intercropping may improve

harvest index and nitrogen use efficiency of crops as well as possibly produce more nutritious food without high inputs.

Table 1.

System	Grain Yield kg ha ⁻¹	Total biomass kg ha ⁻¹ *	Harvest index (HI) *	N % in grain *	N uptake kg ha ⁻¹	NUE kg kg ⁻¹
Monoculture	7383.86	17400	0.44	0.74	177.27	43.60
Intercropped	7094.59	12107	0.59	1.09	159.74	46.61

*statistically significant difference obtained from ANOVA, $p < 0.001$

Acknowledgments

Authors would like to thank Croatian Science Foundation for funding this research through project UIP-7103 "Intercropping of wood species and agricultural crops as an innovative approach in agroecosystems"

References

- Artru, S., Garre, S., Dupraz, C., Hiel, M.-P., Blitz-Frayret, C., & Lassois, L. (2016). Impact of spatio-temporal shade dynamics on wheat growth and yield, perspectives for temperate agroforestry. *European Journal of Agronomy*, 82. <https://doi.org/10.1016/j.eja.2016.10.004>.
- Cannell, M. G. R., Van Noordwijk, M., & Ong, C. K. (1996). The central agroforestry hypothesis: The trees must acquire resources that the crop would not otherwise acquire. *Agroforestry Systems*, 34(1), 27–31. <https://doi.org/10.1007/BF00129630>.
- Dufour L., Metay A., Talbot G., Dupraz C. (2013). Assessing light competition for cereal production in temperate agroforestry systems using experimentation and crop modelling. *Journal of Agronomy and Crop Science*, 199 (3):217–227.
- Gosme, M., Dufour, L., Inurreta-Aguirre, H., & Dupraz, C. (2016). Microclimatic effect of agroforestry on diurnal temperature cycle. 3rd European Agroforestry Conference (May), 183-186.



EURAF2022



Organizing Secretariat
Be tools

Cagliari, Sardinia (Italy)
www.betools.it
office@betools.it

ISBN: 9788897666189